

Properties of Hurwitz Equivalence in the Braid Group of Order n

T. Ben-Itzhak and M. Teicher ¹

ABSTRACT. In this paper we prove certain Hurwitz equivalence properties in B_n . Our main result is that every two Artin's factorizations of Δ_n^2 of the form $H_{i_1} \cdots H_{i_{n(n-1)}}, F_{j_1} \cdots F_{j_{n(n-1)}}$ (with $i_k, j_k \in \{1, \dots, n-1\}$), where $\{H_1, \dots, H_{n-1}\}, \{F_1, \dots, F_{n-1}\}$ are frames, are Hurwitz equivalent. This theorem is a generalization of the theorem we have proved in [6], using an algebraic approach unlike the proof in [6] which is geometric.

The application of the result is applied to the classification of algebraic surfaces up to deformation. It is already known that there exist surfaces that are diffeomorphic to each other but are not deformations of each other (Manetti example). We construct a new invariant based on a Hurwitz equivalence class of factorization, to distinguish among diffeomorphic surfaces which are not deformations of each other. The precise definition of the new invariant can be found in [4] or [5]. The main result of this paper will help us to compute the new invariant.

1 Topological Background

In this section we recall some basic definitions and statements from [3]: Let D be a closed disk on $\mathbb{R}^2$, $K \subset D$ finite set, $u \in \partial D$. Any diffeomorphism of D which fixes K and is the identity on ∂D acts naturally on $\Pi_1 = \Pi_1(D - K, u)$. We say that two such diffeomorphisms of D (which fix K and equal identity on ∂D) are equivalent if they define the same automorphism on $\Pi_1(D - K, u)$. This equivalence relation is compatible with

¹Partially supported by the Emmy Noether Research Institute for Mathematics and the Minerva Foundation of Germany and to the Excellency Center "Group Theoretic Methods in the Study of Algebraic Varieties" of the Israel Science Foundation.

composition of diffeomorphism and thus the equivalence classes form a group.

Definition 1.1. *Braid Group $B_n[D, K]$.*

Let D, K be as above, $n = \#K$, and let $\mathcal{B}$ be the group of all diffeomorphisms β of D such that $\beta(K) = K$, $\beta|_{\partial D} = Id_{\partial D}$. For $\beta_1, \beta_2 \in \mathcal{B}$ we say that β_1 is equivalent to β_2 if β_1 and β_2 define the same automorphism of $\Pi_1(D - K, u)$. The quotient of $\mathcal{B}$ by this equivalence relation is called the braid group $B_n[D, K]$.

Equivalently, if we take the canonical homomorphism $\psi : \mathcal{B} \rightarrow Aut(\Pi_1(D - K, u))$, then $B_n[D, K] = Im(\psi)$. The elements of $B_n[D, K]$ are called braids.

Lemma 1.2. *If $K' \subset D'$ in another pair as above with $\#K' = \#K = n$, then $B_n[D', K']$ is isomorphic to $B_n[D, K]$.*

This gives rise to the definition of B_n :

Definition 1.3. $B_n = B_n[D, K]$ for some D, K with $\#K = n$.

Definition 1.4. $l(q)$.

Let c be a small loop equal to the oriented boundary of a small environment V of w_0 chosen so that $q' = q - (V \cap q)$ is a simple path. Then $l(q) = q' \cup c \cup q'^{-1}$. We also use the notation $l(q)$ for the element of $\Pi_1(D - K, u)$ corresponding to $l(q)$.

Definition 1.5. *A Bush.*

Consider in D an ordered set of simple paths $(T_1, \dots, T_n)$ connecting $w_i \in K$ with u such that:

- (1) $\forall i = 1, \dots, n \quad T_i \cap w_j = \emptyset$ if $i \neq j$.
- (2) $\bigcap_{i=1}^n T_i = u$.
- (3) For a small circle $c(u)$ around u , each $u_i' = T_i \cap c(u)$ is a single point and the order in $(u_1', \dots, u_n')$ is consistent with the positive (“counter-clockwise”) orientation of $c(u)$. We say that two such sets $(T_1, \dots, T_n)$ and $(T'_1, \dots, T'_n)$ are equivalent if $\forall i = 1, \dots, n \quad l(T_i) = l(T'_i)$ in $\Pi_1(D - K, u)$. An equivalent class of such sets is called a bush in $(D - K, u)$. The bush represented by $(T_1, \dots, T_n)$ is denoted by $\langle T_1, \dots, T_n \rangle$.

It is well known that $\Pi_1(D - K, u)$ is a free group.

Definition 1.6. *g-base*

A g-base of $\Pi_1(D - K, u)$ is an ordered free base of $\Pi_1(D - K, u)$ which has the form $(l(T_1), \dots, l(T_n))$ where $\langle T_1, \dots, T_n \rangle$ is a bush in $(D - K, u)$.

Definition 1.7. $H(\sigma)$, *half-twist defined by σ* .

Let D and K be defined as above. Let $a, b \in K$, $K_{a,b} = K \setminus \{a, b\}$ and σ be a simple path (without a self intersection) in $D \setminus \partial D$ connecting a with b such that $\sigma \cap K = \{a, b\}$. Choose a small regular neighborhood U of σ such that $K_{a,b} \cap U = \emptyset$, and an orientation preserving diffeomorphism $\psi : \mathbb{R}^2 \rightarrow \mathbb{C}^1$ such that $\psi(\sigma) = [-1, 1] = \{z \in \mathbb{C}^1 | \operatorname{Re}(z) \in [-1, 1], \operatorname{Im}(z) = 0\}$ and $\psi(U) = \{z \in \mathbb{C}^1 | |z| < 2\}$. Let $\alpha(r), r \geq 0$, be a real smooth monotone function such that $\alpha(r) = 1$ for $r \in [0, 3/2]$ and $\alpha(r) = 0$ for $r \geq 2$. Define a diffeomorphism $h : \mathbb{C}^1 \rightarrow \mathbb{C}^1$ as follows: for $z \in \mathbb{C}^1, z = re^{i\phi}$ let $h(z) = re^{i(\phi + \alpha(r)\pi)}$. It is clear that the restriction of h to $\{z \in \mathbb{C}^1 | |z| \leq 3/2\}$ coincides with the 180° positive rotation, and that the restriction to $\{z \in \mathbb{C}^1 | |z| \geq 2\}$ is the identity map. The diffeomorphism $\psi^{-1} \circ h \circ \psi$ induces a braid called half-twist and denoted by $H(\sigma)$

Definition 1.8. *Frame of $B_n[D, K]$*

Let $K = \{a_1, \dots, a_n\}$ and $\sigma_1, \dots, \sigma_{n-1}$ be a system of simple smooth paths in $D - \partial D$ such that σ_i connects a_i with a_{i+1} and $L = \bigcup \sigma_i$ is a simple smooth path. The ordered system of half-twists $(H_1, \dots, H_{n-1})$ defined by $\{\sigma_i\}_{i=1}^{n-1}$ is called a frame of $B_n[D, K]$

Theorem 1.9. *Let $(H_1, \dots, H_{n-1})$ be a frame of $B_n[D, K]$ then, $B_n[D, K]$ is generated by $\{H_i\}_{i=1}^{n-1}$ and the following is a complete list of relations:*

$H_i H_j = H_j H_i$ if $|i - j| > 1$ (*Commutative relation*)

$H_i H_j H_i = H_j H_i H_j$ if $|i - j| = 1$ (*Triple relation*).

Proof can be found for example in [3].

This theorem provides us with Artin's algebraic definition of Braid group.

2 Definition of Hurwitz Moves

Definition 2.1. *Hurwitz move on G^m (R_k, R_k^{-1})*

Let G be a group, $\vec{t} = (t_1, \dots, t_m) \in G^m$. We say that $\vec{s} = (s_1, \dots, s_m) \in G^m$ is obtained from $\vec{t}$ by the Hurwitz move R_k (or $\vec{t}$ is obtained from $\vec{s}$ by the Hurwitz move R_k^{-1}) if

$$\begin{aligned} s_i &= t_i & \text{for } i \neq k, k+1, \\ s_k &= t_k t_{k+1} t_k^{-1}, & s_{k+1} = t_k. \end{aligned}$$

Definition 2.2. *Hurwitz move on factorization*

Let G be a group and $t \in G$. Let $t = t_1 \cdots t_m = s_1 \cdots s_m$ be two factorized expressions of t . We say that $s_1 \cdots s_m$ is obtained from $t_1 \cdots t_m$ by the Hurwitz move R_k if $(s_1, \dots, s_m)$ is obtained from $(t_1, \dots, t_m)$ by the Hurwitz move R_k .

Definition 2.3. *Hurwitz equivalence of factorization*

The factorizations $s_1 \cdots s_m, t_1 \cdots t_m$ are Hurwitz equivalent if they are obtained from each other by a finite sequence of Hurwitz moves. The notation is $t_1 \cdots t_m \stackrel{HE}{\sim} s_1 \cdots s_m$.

Definition 2.4. *Word in B_n*

A word in B_n is a representation of braid as a sequence of the frame elements and their inverses.

3 Hurwitz Equivalence of Factorizations with Frame Elements

We already proved Proposition 3.3 in [6]. We provide here the proof once again since the logic use is basic for understanding the sequel.

We will need a certain result of Garside:

Claim 3.1. (Garside): *Every two positive words (all generators with positive powers) which are equal are transformable into each other through a finite sequence of positive words, such that each word of the sequence is positive and obtained from the preceding one by a direct application of the commutative relation or the triple relation.*

Proof: [1].

The following is obvious, nevertheless we give a short proof for clarification:

Claim 3.2. *G is a group $g_1, g_2 \in G$:*

1. *If $g_1 g_2 = g_2 g_1$ then $g_1 \cdot g_2 \stackrel{HE}{\sim} g_2 \cdot g_1$.*
2. *If $g_1 g_2 g_1 = g_2 g_1 g_2$ then $g_1 \cdot g_2 \cdot g_1 \stackrel{HE}{\sim} g_2 \cdot g_1 \cdot g_2$.*

Proof:

1. $g_1 \cdot g_2 \xrightarrow{R_1} g_1 g_2 g_1^{-1} \cdot g_1 = g_2 \cdot g_1$.
2. $g_1 \cdot g_2 \cdot g_1 \xrightarrow{R_2} g_1 \cdot g_2 g_1 g_2^{-1} \cdot g_2 \xrightarrow{R_1} g_1 g_2 g_1 g_2^{-1} g_1^{-1} \cdot g_1 \cdot g_2$ but $g_1 g_2 g_1 = g_2 g_1 g_2$, so $g_1 g_2 g_1 g_2^{-1} g_1^{-1} = g_2$ and we get $g_1 \cdot g_2 \cdot g_1 \stackrel{HE}{\sim} g_2 \cdot g_1 \cdot g_2$

From Garside and the above, we get the following proposition:

Proposition 3.3. *Let $H_1, \dots, H_{n-1}$ be a set of generators of B_n and $H_{i_1} \dots H_{i_p} = H_{j_1} \dots H_{j_p}$ two positive words (with $i_k, j_k \in \{1, \dots, n-1\}$) then $H_{i_1} \dots H_{i_p} \stackrel{HE}{\sim} H_{j_1} \dots H_{j_p}$.*

Proof:

Applying 3.1 on $H_{i_1} \dots H_{i_p} = H_{j_1} \dots H_{j_p}$, we get a finite sequence of positive words $\{W_r\}_{r=0}^q$ s.t. $W_0 = H_{i_1} \dots H_{i_p}$, $W_q = H_{j_1} \dots H_{j_p}$ and W_{r+1} is obtained from W_r by a single application of the commutative relation or the triple relation.

An application of the triple relation (as in 3.2) is equal to an application of 2 Hurwitz moves R_{t+1}, R_t . An application of the commutative relation is equal to an application of one Hurwitz move (on the commuting elements). Thus, $W_0 \stackrel{HE}{\sim} W_q$

Definition 3.4. $\Delta_n^2 \in B_n[D, K]$

$$\Delta_n^2 = (H_1 \dots H_{n-1})^n \text{ where } \{H_i\}_{i=1}^{n-1} \text{ is a frame.}$$

We apply 3.3 on Δ_n^2 :

Corollary 3.5. *All Δ_n^2 factorizations $H_{i_1} \cdots H_{i_{n(n-1)}}$ $i_k \in \{1, \dots, n-1\}$ are Hurwitz equivalent.*

Notation 3.6. *We use the notation $a[b]$ for conjugating b to a .*

Proposition 3.7. *Let $\{H(\sigma_i)\}_{i=1}^{n-1}$ be the frame which generates $B_n[D, K]$ and H_ζ a half twist. Then $\{H(\sigma_i)[H_\zeta]\}_{i=1}^{n-1}$ is also a frame.*

Proof:

By III.1.0 of [2], $H(\sigma_i)[H_\zeta] = H(H_\zeta(\sigma_i))$, thus conjugating H_ζ to the elements in the frame is the same as operating H_ζ on each of the elements in $(\sigma_1, \dots, \sigma_{n-1})$, and we get $(H_\zeta(\sigma_1), \dots, H_\zeta(\sigma_{n-1}))$ which also defines a frame since:

- $\bigcup \sigma_i$ is a simple smooth path. H_ζ is a diffeomorphism, so $H_\zeta(\bigcup \sigma_i) = \bigcup H_\zeta(\sigma_i)$ is also a simple smooth path.
- if ζ connects a_i with a_j , $H_\zeta(\sigma_r)$ connects a_r with a_{r+1} when $r \neq i-1, i, j-1, j$. $H_\zeta(\sigma_{i-1})$ connects a_{i-1} with a_j , $H_\zeta(\sigma_i)$ connects a_j with a_{i+1} , $H_\zeta(\sigma_{j-1})$ connects a_{j-1} with a_i , $H_\zeta(\sigma_j)$ connects a_i with a_{j+1} . This means that a_{i-1}, a_{i+1} are connected through a_j , and a_{j-1}, a_{j+1} are connected through a_i as shown in Figure 1.

Proposition 3.8. *Let $(X_1, \dots, X_{n-1})$ be the frame which generates $B_n[D, K]$ and Π is the factorization $X_1 \cdots X_{n-1}$, then:*

1. $X_i \cdot \Pi \stackrel{HE}{\sim} \Pi \cdot X_{i-1}$ for $1 < i \leq n-1$.
2. $X_1 \cdot \Pi \cdot \Pi \stackrel{HE}{\sim} \Pi \cdot \Pi \cdot X_{n-1}$.

Proof:

It is enough to show that the words are equal and by 3.3 the factorizations are Hurwitz equivalent.

1. $X_i \Pi = X_i(X_1 \cdots X_{i-1} X_i \cdots X_{n-1}) = X_1 \cdots X_i X_{i-1} X_i \cdots X_{n-1} = X_1 \cdots X_{i-1} X_i X_{i-1} X_{i+1} \cdots X_{n-1} = (X_1 \cdots X_{i-1} X_i X_{i+1} \cdots X_{n-1}) X_{i-1} = \Pi X_{i-1}$.
2. $X_1 \Pi \Pi = X_1 \Pi X_1 \cdots X_{n-2} X_{n-1}$ and from 1. We get that $X_1 \Pi X_1 \cdots X_{n-2} X_{n-1} = X_1 X_2 \cdots X_{n-1} \Pi X_{n-1}$.




Figure 1: $(H_\zeta(\sigma_1), \dots, H_\zeta(\sigma_{n-1}))$.

Proposition 3.9. *Let $\{X_i\}_{i=1}^{n-1}$ be the frame which generates $B_n[D, K]$. For each $1 \leq j \leq n-1$ the Δ_n^2 factorizations $(X_1 \cdots X_{n-1})^n$ and $(X_1[X_j] \cdots X_{n-1}[X_j])^n$ are Hurwitz equivalent.*

Proof:

We first consider the two extreme cases where $j = n-1$ and $j = 1$.

In the case where, $j = n-1$, by performing the series of Hurwitz moves $\{R_{m(n-1)-1}^{-1}\}_{m=1}^n$ on $(X_1 \cdots X_{n-1})^n$ (each of the Hurwitz moves is acting on one of the n factors in the product) we get that

$$(X_1 \cdots X_{n-3} \cdot X_{n-2} \cdot X_{n-1})^n \stackrel{HE}{\rightsquigarrow} (X_1 \cdots X_{n-3} \cdot X_{n-1} \cdot X_{n-2}[X_{n-1}])^n.$$

Since $X_i[X_{n-1}] = X_i$ for $i \neq n-2$, all elements in the factorization are of the form $X_i[X_{n-1}]$ and by 3.7 they are all elements of the same frame. $(X_1 \cdots X_{n-1} \cdot X_{n-2}[X_{n-1}])^n$ and $(X_1 \cdots X_{n-2}[X_{n-1}] \cdot X_{n-1})^n$ are both Δ_n^2 factorizations.

Having the same frame elements and from 3.5 they are Hurwitz equivalent, and we get that:

$$(X_1[X_{n-1}] \cdots X_{n-2}[X_{n-1}] \cdot X_{n-1}[X_{n-1}])^n \stackrel{HE}{\rightsquigarrow} (X_1 \cdots X_{n-1})^n.$$

The case where $j = 1$ is similar by performing the series of Hurwitz moves $\{R_{m(n-1)+1}^{-1}\}_{m=0}^{n-1}$. In the case where $1 < j < n-1$, we perform $\{R_{m(n-1)+j}^{-1}\}_{j=0}^{n-1}$

and get:

$$(X_1, \dots, X_{j-1} \cdot X_j \cdot X_{j+1} \dots X_{n-1})^n \stackrel{HE}{\sim} (X_1, \dots, X_{j-2} \cdot X_j \cdot X_{j-1}[X_j] \cdot X_{j+1} \dots X_{n-1})^n.$$

Since X_j commutes with $X_1, \dots, X_{j-2}$, then by 3.2,

$$(X_1, \dots, X_{j-2} \cdot X_j \cdot X_{j-1}[X_j] \cdot X_{j+1} \dots X_{n-1})^n \stackrel{HE}{\sim} (X_j \cdot X_1, \dots, X_{j-2} \cdot X_{j-1}[X_j] \cdot X_{j+1} \dots X_{n-1})^n.$$

(In fact we performed $\{\{R_{m(n-1)+k}\}_{k=j-2}^1\}_{m=0}^{n-1}$).

Again, X_j commutes with $X_{j+2}, \dots, X_{n-1}$ so by 3.2:

$$(X_j \cdot X_1 \dots X_{j-2} \cdot X_{j-1}[X_j] \cdot X_{j+1} \dots X_{n-1})^n \stackrel{HE}{\sim} (X_j \cdot X_1 \dots X_{j-2} \cdot X_{j-1}[X_j] \cdot X_{j+1} \cdot X_j \cdot X_{j+2} \dots X_{n-1}) \cdot (X_1 \dots X_{j-2} \cdot X_{j-1}[X_j] \cdot X_{j+1} \cdot X_j \cdot X_{j+2} \dots X_{n-1})^{n-2} (X_1 \dots X_{j-2} \cdot X_{j-1}[X_j] \cdot X_{j+1} \cdot X_{j+2} \dots X_{n-1})$$

which is Hurwitz equivalent to

$$(X_j \cdot X_1 \dots X_{j-2} \cdot X_{j-1}[X_j] \cdot X_j \cdot X_{j+1}[X_j] \cdot X_{j+2} \dots X_{n-1}) \cdot (X_1 \dots X_{j-2} \cdot X_{j-1}[X_j] \cdot X_j \cdot X_{j+1}[X_j] \cdot X_{j+2} \dots X_{n-1})^{n-2} (X_1 \dots X_{j-2} \cdot X_{j-1}[X_j] \cdot X_{j+1} \cdot X_{j+2} \dots X_{n-1})$$

(In fact we performed $\{R_{j+m(n-1)+1}^{-1}\}_{m=0}^{n-2}$)

Let $(A_1, \dots, A_{n-1})$ be the conjugated frame s.t. $A_i = X_i[X_j]$ and $\Pi = A_1 \dots A_{n-1}$.

The previous factorization can be written as

$$A_j \cdot \Pi^{n-1} \cdot A_1 \dots A_{j-1} \cdot X_{j+1} \cdot A_{j+2} \dots A_{n-1}.$$

by 3.8 (1) this factorization is Hurwitz equivalent to:

$$\Pi^{j-1} \cdot A_1 \cdot \Pi^{n-j} \cdot A_1 \dots A_{j-1} \cdot X_{j+1} \cdot A_{j+2} \dots A_{n-1}.$$

and by 3.8 (2) we get that

$$\Pi^{j-1} \cdot A_1 \cdot \Pi^{n-j} \cdot A_1 \dots A_{j-1} \cdot X_{j+1} \cdot A_{j+2} \dots A_{n-1}$$

is Hurwitz equivalent to

$$\Pi^{j+1} \cdot A_{n-1} \cdot \Pi^{n-j-2} \cdot A_1 \cdots A_{j-1} \cdot X_{j+1} \cdot A_{j+2} \cdots A_{n-1}$$

Using 3.8 (1) $n - j - 2$ times we get that the above is Hurwitz equivalent to

$$\Pi^{n-1} \cdot A_{j+1} \cdot A_1 \cdots A_{j-1} \cdot X_{j+1} \cdot A_{j+2} \cdots A_{n-1}.$$

Since A_{j+1} commutes with $A_1, \dots, A_{j-1}$ then by 3.2:

$$\begin{aligned} & \Pi^{n-1} \cdot A_{j+1} \cdot A_1 \cdots A_{j-1} \cdot X_{j+1} \cdot A_{j+2} \cdots A_{n-1} \stackrel{HE}{\sim} \\ & \Pi^{n-1} \cdot A_1 \cdots A_{j-1} \cdot A_{j+1} \cdot X_{j+1} \cdot A_{j+2} \cdots A_{n-1}. \end{aligned}$$

By performing $R_{(n-1)^2+j}$ (on A_{j+1}), the factorization is Hurwitz equivalent to:

$$\Pi^{n-1} \cdot A_1 \cdots A_{j-1} \cdot X_{j+1} [(X_{j+1}[X_j])^{-1}] \cdot A_{j+1} \cdot A_{j+2} \cdots A_{n-1}$$

which is exactly $(A_1 \cdots A_{n-1})^n$.

Proposition 3.10. *Let $b \in B_n[D, K]$, then the factorizations $(X_1 \cdots X_{n-1})^n$ and $(X_1[b] \cdots X_{n-1}[b])^n$ are Hurwitz equivalent.*

Proof:

From [1], any braid $P \in B_n[D, K]$ can be represented as $P = \Delta_n^r \overline{P}$ where $\overline{P}$ is a positive braid. If $r \geq 0$ then clearly $A[P]$ is a conjugation of a positive braid. Assume $r < 0$, take m s.t. $2m > -r$ and since Δ_n^2 is the generator of the center of $B_n[D, K]$, then $A[P] = A[\Delta_n^r \overline{P}] = A[\Delta_n^{r+2m} \overline{P}]$ which is a conjugation of a positive braid.

For the rest of the proof we may assume that P is a positive braid.

The rest of the proof will be performed by induction on the length of the positive braid P . The case where $P = X_{i_1}$ was already proven.

If $P = X_{i_1} \cdots X_{i_d} X_{i_{d+1}}$, by the induction assumption,

$$(X_1 \cdots X_{n-1})^n \stackrel{HE}{\sim} (X_1[X_{i_d} \cdots X_{i_1}] \cdots X_{n-1}[X_{i_d} \cdots X_{i_1}])^n.$$

The new elements are also a frame, so by applying 3.9 we get the equivalent factorization:

$$((X_1[X_{i_d} \cdots X_{i_1}])[X_{i_{d+1}}[X_{i_d} \cdots X_{i_1}]] \cdots (X_{n-1}[X_{i_d} \cdots X_{i_1}])[X_{i_{d+1}}[X_{i_d} \cdots X_{i_1}]])^n.$$

For each $1 \leq j \leq n-1$, $(X_j[X_{i_d} \dots X_{i_1}])[X_{i_{d+1}}[X_{i_d} \dots X_{i_1}]] = X_j[X_{i_{d+1}} \dots X_{i_1}]$ and therefore,

$$(X_1 \cdots X_{n-1})^n \stackrel{HE}{\sim} (X_1[X_{i_{d+1}} \dots X_{i_1}] \cdots X_{n-1}[X_{i_{d+1}} \dots X_{i_1}])^n.$$

□

4 The Main Result

In the following section we will apply the previous proposition on all frames, by proving that every two frames of $B_n[D, K]$ are conjugated.




Figure 2: g-base of a frame.

Definition 4.1. *g-base of a frame F .*

Let $F = (F_1, \dots, F_{n-1})$ ($F_i = H(\sigma_i)$) be a frame of $B_n[D, K]$. Let the system of paths $(T_1, \dots, T_n)$, T_i connecting u with $a_i \in K$ create a bush s.t.

$$T_i \cap \sigma_j = \begin{cases} \phi, & \text{if } j \neq i, i-1 \\ a_i, & \text{if } j = i \text{ or } j = i-1 \end{cases}.$$

Let $\gamma_i = l(T_i)$ then $\Gamma = (\gamma_1, \dots, \gamma_n)$ is the g-base of the frame F as shown in Figure 2.

Proposition 4.2. *If $\Gamma = (\gamma_1, \dots, \gamma_n)$ is the g-base of two frames, then the frames are equal.*

Proof:

Assuming that Γ is the g-base of the two frames F_1 and F_2 where , $F_1 = (H(\sigma_1), \dots, H(\sigma_{n-1}))$ and $F_2 = (H(\zeta_1), \dots, H(\zeta_{n-1}))$ and $F_1 \neq F_2$. Since $F_1 \neq F_2$ then $\exists i \in \{1, \dots, n-1\}$ s.t. $H(\sigma_i) \neq H(\zeta_i)$.

Γ is the g-base of both frames, then σ_i and ζ_i connecting a_i with a_{i+1} . $H(\sigma_i) \neq H(\zeta_i)$, therefore, ζ_i and σ_i are “separated” by a point $a_j \neq a_i, a_{i+1}$ as shown in Figure 3. The two paths start at a_i and end at a_{i+1} therefore creating a cycle with a_j inside. Since T_j connects a_j with u , $T_j \cap \sigma_i \neq \emptyset$ or $T_j \cap \zeta_i \neq \emptyset$ and therefore Γ is not the g-base of both F_1 and F_2 .




Figure 3: Two frames with the same g-base.

Proposition 4.3. *Let $\Gamma = (\gamma_1, \dots, \gamma_n)$ be the g-base of the frame $F = (H(\sigma_1), \dots, H(\sigma_{n-1}))$ and let H_ζ be a half twist. Then $(H_\zeta(\gamma_1), \dots, H_\zeta(\gamma_n))$ is the g-base of the frame $(H(\sigma_1)[H_\zeta], \dots, H(\sigma_{n-1})[H_\zeta])$.*

Proof:

It was already proven in [3] that any braid $b \in B_n[D, K]$ transforms a g-base to a g-base. By Proposition 3.7, $(H_\zeta(\sigma_1), \dots, H_\zeta(\sigma_{n-1}))$ defines a frame.

Since H_ζ is a diffeomorphism, acting on $\sigma_1, \dots, \sigma_{n-1}, \gamma_1, \dots, \gamma_n$ does not create new intersection points while swapping a_i with a_j for both frame and g-base elements. Therefore, $H_\zeta(\Gamma)$ is the g-base of the conjugated frame.

Corollary 4.4. *Let $\Gamma = (\gamma_1, \dots, \gamma_n)$ be the g-base of $F = (H(\sigma_1), \dots, H(\sigma_{n-1}))$ and $b \in B_n[D, K]$, then $(b(\gamma_1), \dots, b(\gamma_n))$ is the g-base of the frame $(H(\sigma_1)[b], \dots, H(\sigma_{n-1})[b])$.*

Proof:

Induction on the length of P using Proposition 4.3.

Proposition 4.5. *Let $(H(\sigma_1), \dots, H(\sigma_{n-1}))$ be a frame, then for every frame $(H(\sigma'_1), \dots, H(\sigma'_{n-1}))$ there exist a braid $b \in B_n[D, K]$ s.t. $(H(\sigma_1)[b], \dots, H(\sigma_{n-1})[b]) = (H(\sigma'_1), \dots, H(\sigma'_{n-1}))$.*

Proof:

Let Γ and Γ' be the g-bases of $(H(\sigma_1), \dots, H(\sigma_{n-1}))$ and $(H(\sigma'_1), \dots, H(\sigma'_{n-1}))$. In [3] it was proved that there exists a unique braid $b \in B_n[D, K]$ s.t. $b(\Gamma) = \Gamma'$. From Corollary 4.4 $(H(\sigma_1)[b], \dots, H(\sigma_{n-1})[b])$ is a frame with a g-base Γ' .

From Proposition 4.2, Γ' has only one frame, therefore, $(H(\sigma_1)[b], \dots, H(\sigma_{n-1})[b]) = (H(\sigma'_1), \dots, H(\sigma'_{n-1}))$

We now come to the main result:

Theorem 4.6. *Let $(H_1, \dots, H_{n-1})$ and $(F_1, \dots, F_{n-1})$ be two frames of $B_n[D, K]$, then for every two braids s.t. $\Delta_n^2 = F_{i_1} \dots F_{i_{n(n-1)}} = H_{j_1} \dots H_{j_{n(n-1)}}$ the factorizations $F_{i_1} \dots F_{i_{n(n-1)}}$ and $H_{j_1} \dots H_{j_{n(n-1)}}$ are Hurwitz equivalent.*

Proof:

From Corollary 3.5, $F_{i_1} \dots F_{i_{n(n-1)}} \stackrel{HE}{\sim} (F_1 \dots F_{n-1})^n$ and $H_{j_1} \dots H_{j_{n(n-1)}} \stackrel{HE}{\sim} (H_1 \dots H_{n-1})^n$.

From Proposition 4.5, $\exists b \in B_n[D, K]$ s.t. $(F_1, \dots, F_{n-1}) = (H_1[b], \dots, H_{n-1}[b])$, so from Proposition 3.10, $(F_1 \dots F_{n-1})^n \stackrel{HE}{\sim} (H_1 \dots H_{n-1})^n$ and therefore, $F_{i_1} \dots F_{i_{n(n-1)}}, H_{j_1} \dots H_{j_{n(n-1)}}$ are Hurwitz equivalent. $\square$

References

- [1] F. A. Garside, *The Braid group and other groups* Quart. J. Math Oxford (2) 78 (1969), 235-254.
- [2] B. Moishezon and M. Teicher, *Braid group technique in complex geometry V: The fundamental group of a complement of a branch curve of a veronese generic projection*, Communication in Analysis and Geometry 4, No. 11, (1996), 1-120
- [3] B. Moishezon and M. Teicher, *Braid group techniques in complex geometry I, Line arrangements in $\mathbb{CP}^2$* , Contemporary Math. 78 (1988), 425-555.
- [4] Vik. S. Kulikov and M. Teicher, *Braid monodromy factorization and diffeomorphism types*, Izvestia, Journal of the Russian Academy of Science, Tom 64, 2, 2000, 89-120.
- [5] M. Teicher, *Braid monodromy type of 4 manifolds*, Preprint.
- [6] M. Teicher and T. Ben-Itzhak, *Hurwitz equivalence in braid group B_3* , Preprint.